

DTIC FILE COPY

AD-A227 338

A Presentation on Explicit Normal Modes for a Perturbed Ocean Model By Perturbation

**A Paper Presented at the
First IMACS Symposium on
Computational Acoustics,
6-8 August 1986, New Haven, Connecticut**

Mark D. Duston
Robert P. Gilbert
Ghasi R. Verma
David H. Wood
Surface Ship Sonar Department



**Naval Underwater Systems Center
Newport, Rhode Island / New London, Connecticut**

Approved for public release; distribution limited.


DTIC
ELECTE
OCT 12 1990
S E D
cc

90 10 12 002

Preface

The work reported in this document was completed under NUSC Project No. A92045, Dr. Rolf Kasper, Principal Investigator. A portion of this research was done while one of us (Gilbert) was on an Intergovernmental Personnel Act Mobility Assignment to the Naval Underwater Systems Center from the University of Delaware. Another portion of this research was completed while two of us (Duston and Verma) participated in the U.S. Navy-ASEE Summer Faculty Research Program at the Naval Underwater Systems Center, New London Laboratory, in conjunction with NUSC Independent Research Project A92045.

Reviewed and Approved: 7 October 1986

A handwritten signature in cursive script, reading "W. A. Von Winkle".

W. A. Von Winkle
Associate Technical Director
for Technology

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) TD 7791			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Naval Underwater Systems Center		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code). New London Laboratory New London, CT 20362			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) A PRESENTATION ON EXPLICIT NORMAL MODES FOR A PERTURBED OCEAN MODEL BY PERTURBATION					
12. PERSONAL AUTHOR(S) Mark D. Duston, Robert P. Gilbert, Ghasi R. Vermi and David H. Wood					
13a. TYPE OF REPORT		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 1986 October 7	
				15. PAGE COUNT	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Wave Propagation Perturbation		
			Transmutation Computer Modeling		
			Acoustics		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>This document is based on a presentation at the First IMACS Symposium on Computational Acoustics, held in New Haven, Connecticut, 6-8 August 1986. (The title was changed from "Effects of Matching Boundary Conditions on Normal Modes Using Perturbation Theory.")</p> <p>Subject to symplifying assumptions, we are able to use the method of normal modes, involving the eigenvalues and eigenfunctions of a depth dependent Sturm-Liouville problem. We allow the constant sound speed of the idealized uniform ocean model to be modified by a function that is multiplied by a small parameter, epsilon. We investigate the changes that result, using both the classical theory of perturbation for Sturm-Liouville problems and a newer <u>transmutation</u> approach. We are able to use the transmutation approach in a way</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL David H. Wood			22b. TELEPHONE (Include Area Code) (203) 440-4831		22c. OFFICE SYMBOL Code 3332

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete.SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

19. ABSTRACT (Cont'd.)

that overcomes its main disadvantage, at least to first order in epsilon.

- We then obtain explicit formulas for both the eigenvalues and eigenfunctions to first order, without having to expand the eigenfunctions in an infinite series, as is done in the classical approach. Finally, we show that the new formulas are consistent with the classical infinite series.



EXPLICIT NORMAL MODES FOR A PERTURBED OCEAN MODEL BY TRANSMUTATION

MARK D. DUSTON
ROBERT P. GILBERT
GHASI R. VERMA
DAVID H. WOOD

Graph 1


The theoretical foundations of normal mode theory and its use are already well established. In the future we wish to consider range dependent problems in terms of partitioning, so that the range dependent effects will be corrected at the boundaries of the partitions and the solutions allowed to propagate further. In order to apply this approach it is necessary to find the normal modes at each boundary. We are interested, therefore, in finding a method that will allow us to calculate the normal modes in a faster, more efficient manner. One of the methods that holds promise of attaining this objective is that of transmutation theory. Transmutation theory is essentially a generalization of the method of integral transformation.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	





THE IDEALIZED OCEAN



$$\left(p_{rr}(r,z) + \frac{1}{r} p_r(r,z) + p_{zz}(r,z) + k^2 p(r,z) = 0, \right. \\ \left. \text{WITH } p(r,0) = 0 \text{ AND } p_z(r,h) = 0. \right)$$

WE SOLVE USING SEPARATION OF VARIABLES,

$$p(r,z) = \phi(z) \theta(r).$$

Graph 2

We are dealing at this point with a fairly simplified model of the ocean and make the following assumptions:

- 1) uniform depth,
- 2) ocean is an isotropic medium,
- 3) sound speed a constant
- 4) pressure $p=0$ at the surface, and
- 5) ocean and its bottom are eventually (at some depth h) underlaid by a rigid surface.

Under these assumptions the excess pressure p satisfies the Helmholtz Equation.

$$p_{rr}(r,z) + \frac{1}{r} p_r(r,z) + p_{zz}(r,z) + k^2 p(r,z) = 0,$$

with the boundary conditions

$$p(r,0) = 0 \quad \text{and} \quad p_z(r,h) = 0.$$

The second boundary condition (the boundary condition at h) is that of a rigid subbottom. We solve in a standard manner by using the method of separation of variables $p(r,z) = \psi(z)\theta(r)$.



THE IDEALIZED STURM-LIOUVILLE PROBLEM

THE IDEALIZED NORMAL MODES SATISFY

$$\phi''(z) + k^2 \phi(z) = \ell \phi(z),$$

$$\phi(0) = 0 \quad \text{AND} \quad \phi'(h) = 0$$

WITH L_2 NORMALIZATION THIS HAS SOLUTIONS

$$\phi_m(z) = \sqrt{\frac{h}{2}} \sin \frac{(2m-1)\pi z}{2h}$$

$$m = 1, 2, 3, \dots$$

$$\ell_m = k^2 - \left[\frac{(2m-1)\pi}{2h} \right]^2$$

Graph 3

It is well known that for the idealized ocean with constant sound speed the normal modes satisfy an idealized Sturm-Liouville problem given by

$$\phi''(z) + k^2 \phi(z) = \ell \phi(z),$$

and boundary conditions

$$\phi(0) = 0 \quad \text{and} \quad \phi'(h) = 0.$$

It is also well known that this problem has a complete set of eigenfunctions given by

$$\phi_m(z) = \sqrt{\frac{h}{2}} \sin \frac{(2m-1)\pi z}{2h} \quad m = 1, 2, \dots$$


and corresponding discrete eigenvalues given by

$$\ell_m = k^2 - \left[\frac{(2m-1)\pi}{2h} \right]^2 \quad m = 1, 2, \dots$$

These eigenfunctions have been L_2 normalized.



THE PERTURBED STURM-LIOUVILLE PROBLEM



IN AN OCEAN WHERE THE SOUND SPEED IS A FUNCTION OF DEPTH ONLY THE NORMAL MODES SATISFY

$$\psi''(z) + k^2 n^2(z) \psi(z) = \lambda \psi$$

WITH THE CONDITIONS

$$\psi(0) = 0 \text{ AND } \psi'(h) = 0.$$

Graph 4

We wish to solve a similar problem associated with amore general ocean model. The condition that the sound speed be constant is replaced with the assumption that the sound speed is a function of the depth only, the other assumptions remain unchanged. We consider this model to be the perturbed ocean model. The depth dependent equation gives a perturbed Sturm-Liouville problem for the normal modes which satisfy

$$\psi''(z) + k^2 n^2(z) \psi(z) = \lambda \psi(z)$$

with the boundary conditions expressed as

$$\psi(0) = 0 \quad \text{and} \quad \psi'(h) = 0.$$



THE INDEX OF REFRACTION

$$n^2(z) = 1 + \epsilon s(z)$$

CONTAINS A PERTURBATION $\epsilon s(z)$.

THE CASE $\epsilon = 0$ REPRESENTS AN IDEALIZED OCEAN
WITH CONSTANT SOUND SPEED.

Graph 5

The quantity $n^2(z)$ is the index of refraction and we represent it as

$$n^2(z) = 1 + \epsilon s(z)$$

where the quantity $\epsilon s(z)$ is considered the perturbation. We introduce the parameter ϵ which reflects the size of the perturbation from its average value. We assume that the function $s(z)$ is known for the case of interest. When $\epsilon=0$ we have recovered the idealized ocean with constant sound speed.



THE GENERAL PERTURBATION APPROACH

FIND THE EIGENFUNCTIONS ψ_n AND THE EIGENVALUES λ_n OF THE PERTURBED PROBLEM IN TERMS OF THE EIGENFUNCTIONS ϕ_n AND EIGENVALUES ℓ_n OF THE IDEALIZED PROBLEM AND THE PERTURBATION $\epsilon s(z)$.

Graph 6

In a perturbation approach we find the eigenfunctions ψ_m and eigenvalues λ_m of the perturbed (depth dependent) in terms of the eigenfunctions ϕ_m and eigenvalues ℓ_m of an idealized problem (which we know) and the perturbation $\epsilon s(z)$ (which we also know). Specifically we look for the changes or corrections which must be made to the idealized eigenfunctions and eigenvalues.



THE CLASSICAL APPROACH

EXPAND IN POWER SERIES IN ϵ

$$\lambda_m = \ell_m + \epsilon \lambda_m^{(1)} + \epsilon^2 \lambda_m^{(2)} + \dots$$

AND

$$\psi_m(z) = \phi_m(z) + \epsilon \psi_m^{(1)}(z) + \epsilon^2 \psi_m^{(2)}(z) + \dots,$$

BUT OBTAIN EACH TERM OF $\psi_m(z)$ ONLY AS A FOURIER SERIES

$$\psi_m^{(i)}(z) = \sum_{p=1}^{\infty} \alpha_{mp}^{(i)} \phi_p(z).$$

Graph 7

The first approach we examine is the classical perturbation approach found in Titchmarsh. The eigenvalues and eigenfunctions of the perturbed problem are expanded in power series of the parameter ϵ .

The perturbed eigenvalue equals the idealized eigenvalue plus corrections,

$$\lambda_m = \ell_m + \epsilon \lambda^{(1)} + \epsilon^2 \lambda^{(2)} + \dots$$

The perturbed eigenfunctions equals the idealized eigenfunctions plus corrections,

$$\psi_m(z) = \phi_m(z) + \epsilon \psi_m^{(1)}(z) + \epsilon^2 \psi_m^{(2)}(z) + \dots$$

However, we find the corrections to the perturbed eigenfunctions only as a Fourier series of the idealized eigenfunctions and we must find the Fourier coefficients α in terms of the idealized eigenfunction, idealized eigenvalues and the perturbation.



THE CLASSICAL RESULTS

THE CORRECTIONS TO FIRST ORDER IN ϵ ARE

$$\lambda_m^{(1)} = \frac{2k^2}{h} \int_0^h s(z) \left[\sin \frac{(2m-1)\pi z}{2h} \right]^2 dz.$$

THE FOURIER COEFFICIENTS ARE

$$\alpha_{mn}^{(1)} = \frac{2k^2}{h} \int_0^h s(z) \sin \frac{(2m-1)\pi z}{2h} \sin \frac{(2n-1)\pi z}{2h} dz,$$

AND

$$\alpha_{mm}^{(1)} = 0.$$

Graph 8

In a fairly straightforward manner the explicit equations for the corrections may be derived. We exhibit the corrections to the first order in ϵ . The first order correction to the m^{th} eigenvalue is given by the formula

$$\lambda^{(1)} = \frac{2k^2}{h} \int_0^h s(z) \left[\sin \frac{(2m-1)\pi z}{2h} \right]^2 dz.$$

The first order correction to the eigenfunctions is given in terms of an infinite fourier series whose coefficients are given by

$$\alpha_{mn}^{(1)} = \frac{2k^2}{h} \int_0^h s(z) \sin \frac{(2m-1)\pi z}{2h} \sin \frac{(2n-1)\pi z}{2h} dz \quad \text{and} \quad \alpha_{mm}^{(1)} = 0.$$



THE TRANSMUTATION APPROACH

WE WANT TO TRANSMUTE SOLUTIONS OF AN IDEALIZED STURM-LIOUVILLE PROBLEM

$$\phi''(z) + k^2 \phi(z) = \lambda \phi(z)$$

INTO SOLUTIONS OF THE PERTURBED STURM-LIOUVILLE PROBLEM

$$\psi''(z) + k^2 n^2(z) \psi(z) = \lambda \psi(z).$$

Graph 9

We say we wish to transmute the solutions of an idealized Sturm-Liouville problem into the solutions of the perturbed Sturm-Liouville problem. A transmutation refers to a generalization of the method of integral transformation. It has the advantage that the choice of the idealized system (the system for which the solutions are known) is at our discretion, we have only to find the proper kernel function for the integral transform. As a rule of thumb one chooses the idealized problem to be as similar as possible to the perturbed problem. We choose for our idealized problem a Sturm-Liouville problem with

$$\phi''(z) + k^2 \phi(z) = \lambda \phi(z)$$

and wish to transmute the solutions into solutions of the perturbed Sturm-Liouville problem with

$$\psi''(z) + k^2 n^2(z) \psi(z) = \lambda \psi(z).$$



THE TRANSMUTATION OPERATION COMPUTES

$$\psi(z) = \varphi(z) + \int_h^z K(z,s) \varphi(s) ds$$

IN TERMS OF THE TRANSMUTATION KERNEL $K(z,s)$
WHICH MUST SATISFY

$$\frac{\partial^2 K}{\partial z^2} - \frac{\partial^2 K}{\partial s^2} + k^2 (n^2(z) - 1) K = 0.$$

Graph 10

We use a transform of the type given by

$$\psi_m(z) = \phi_n(z) + \int_h^z K(z,s) \phi(s) ds$$

where we must specify the kernel function $K(z,s)$.

where the kernel function $K(z,s)$ must be determined. We substitute the transmutation into the separated ordinary differential equation for the variable coefficient Helmholtz equation. This yields a partial differential equation for the kernel function $K(z,s)$

$$\frac{\partial^2 K}{\partial z^2} - \frac{\partial^2 K}{\partial s^2} + k^2 \epsilon_s(z) K = 0.$$



TO SATISFY THE BOUNDARY CONDITION

$$\psi'(h) = 0,$$


WE ASK THAT THE
TRANSMUTATION PRESERVE
THE BOUNDARY CONDITION
AT THE BOTTOM,

$$\phi'(h) = 0, \text{ WHICH GIVES}$$

$$\frac{\partial}{\partial s} K(z, h) = 0,$$

AND

$$2 \frac{\partial}{\partial z} K(z, z) + k^2 (n^2(z) - 1) = 0.$$



Graph 11

We now invoke our boundary conditions for the Sturm-Liouville problem with variable coefficients. We ask that $\psi(0)=0$ and $\psi'(h)=0$. We impose the same bottom boundary condition $\phi'(h)=0$ on the solutions of the Sturm-Liouville problem with constant coefficients. Combining this with the given transmutation we obtain two additional constraints on the kernel function $K(z, s)$

$$\frac{\partial}{\partial s} K(z, h) = 0 \quad \text{and} \quad 2 \frac{\partial}{\partial z} K(z, z) + k^2 \epsilon s(z) = 0.$$

These two conditions and the PDE for $K(z, s)$ are sufficient to uniquely determine the kernel function

It would seem that we have made the problem more difficult, we had reduced the problem to an ODE and now we must solve a PDE. Why not directly solve the original Helmholtz equation? The answer is that we will not directly solve the problem but instead will show a way in which the kernel function $K(z, s)$ may be approximated as a power series in the parameter ϵ .



ONE METHOD OF SOLVING FOR THE KERNEL, $K(z,s)$:

LET $M(\xi, \eta) = K(z,s)$ WHERE $2\xi = z + s - 2h$, $2\eta = z - s$.

THEN

$$2 M_{\xi\eta} + k^2(n^2(\xi + \eta + h) - 1) M = 0.$$

EXPAND $M(\xi, \eta)$ AS

$$M(\xi, \eta) = \sum_{p=1}^{\infty} k^{2p} M^{(p)}(\xi, \eta), \text{ WHERE}$$

$$2M^{(1)}(\xi, \eta) = - \int_0^{\xi} (n^2(\alpha+h) - 1) d\alpha - \int_0^{\eta} (n^2(\beta+h) - 1) d\beta$$

AND

$$M^{(p+1)}(\xi, \eta) = - \int_0^{\eta} \int_0^{\xi} M^{(p)}(\alpha, \beta) (n^2(\alpha + \beta + h) - 1) d\alpha d\beta.$$

Graph 12

One method of solving for the kernel function $K(z,s) = M(\xi, \eta)$ is to transform into characteristic coordinates given by

$$2\xi = z + s - 2h \text{ and } 2\eta = z - s.$$

We then get the PDE for $M(\xi, \eta)$ which is

$$2 M_{\xi\eta} + k^2 n^2(\xi + \eta + h) M = 0.$$

We next do a Born expansion of M in powers of the parameter k^2 . This yields a sequence of integrals where the first coefficient of the expansion is given by

$$2 M^{(1)}(\xi, \eta) = - \int_0^{\xi} \epsilon s(\alpha+h) d\alpha - \int_0^{\eta} \epsilon s(\beta+h) d\beta,$$

and successive coefficients in the expansion are given by

$$M^{(p+1)}(\xi, \eta) = - \int_0^{\eta} \int_0^{\xi} M^{(p)}(\alpha, \beta) \epsilon s(\alpha+\beta+h) d\alpha d\beta.$$



INVERSION OF COORDINATES IN $M(\xi, \eta)$ GIVES

$$K(z, s) = \sum_{p=1}^{\infty} \epsilon^p K^{(p)}(z, s)$$

$$2\epsilon K^{(1)}(z, s) = - \int_h^{\frac{z+s}{2}} k^2(n^2(\alpha) - 1) d\alpha - \int_h^{\frac{z-s+2h}{2}} k^2(n^2(\alpha) - 1) d\alpha$$

AND

$$\epsilon K^{(p+1)}(z, s) = - \int_0^{\frac{z-s}{2}} \int_0^{\frac{z+s-2h}{2}} K^{(p)}(\alpha, \beta) k^2(n^2(\alpha + \beta + h) - 1) d\alpha d\beta.$$

Graph 13

Inverting coordinates from (ξ, η) back into (z, s) we obtain an expansion of the kernel function $K(z, s)$. The first term of the the expansion is

$$2\epsilon K^{(1)}(z, s) = -k^2 \int_h^{(z+s)/2} \epsilon s(\alpha) d\alpha - k^2 \int_h^{(z-s+2h)/2} \epsilon s(\alpha) d\alpha$$

and the successive terms of the expansion are given by

$$\epsilon K^{(p+1)}(z, s) = -k^2 \int_0^{(z-s)/2} \int_0^{(z+s-2h)/2} K^{(p)}(\alpha, \beta) \epsilon s(\alpha + \beta + h) d\alpha d\beta.$$

We have now obtained explicit formulas for the power series expansion of the kernel $K(z, s)$ in the parameter ϵ .



TO SATISFY THE BOUNDARY CONDITION AT THE SURFACE $\phi(s)$ IS CONSTRAINED BY

$$0 = \psi(0) = \phi(0) + \int_h^0 K(0,s) \phi(s) ds.$$

THEREFORE WE DEFINE

$$\theta(z) = \phi(z) - \phi(0).$$

NOTICE THAT

$$\theta''(z) + k^2 (\theta(z) + \phi(0)) = \lambda(\theta(z) + \phi(0));$$

$$\theta(0) = 0 \text{ AND } \theta'(h) = 0.$$

ITS EXPANSION IS

$$\theta(z) = \theta^{(0)}(z) + \epsilon \theta^{(1)}(z) + \epsilon^2 \theta^{(2)}(z) + \dots.$$

Graph 14

We exhibit the value of the transmuted solution at the surface and see that, in general the transmutation does not preserve the surface boundary condition. Therefore we construct a new function

$$\theta(z) = \phi(z) - \phi(0)$$

and see that this function satisfies the same boundary conditions as $\psi(z)$. We then expand this new function in power series in ϵ .



**SUBSTITUTING THE EXPANSION OF $\theta(z)$ INTO
THE STURM-LIOUVILLE PROBLEM GIVES**

ORDER ϵ^0 :

$$\theta^{(0)'''} + (k^2 - \lambda^{(0)}) \theta^{(0)} = 0$$

ORDER ϵ :

$$\theta^{(1)'''} + (k^2 - \lambda^{(0)}) \theta^{(1)} - \lambda^{(1)} \theta^{(0)} = (\lambda^{(0)} - k^2) \phi^{(1)}(0)$$

ORDER ϵ^2 :

$$\theta^{(2)'''} + (k^2 + \lambda^{(0)}) \theta^{(2)} - \lambda^{(1)} \theta^{(1)} - \lambda^{(2)} \theta^{(0)} = (\lambda^{(0)} - k^2) \phi^{(2)}(0) + \lambda^{(1)} \phi^{(1)}(0).$$

Graph 15

Substituting $\theta(z)$ into the idealized Sturm-Liouville problem with constant coefficients and isolating the terms with corresponding powers of ϵ we obtain a sequence of ODE's each with the same boundary conditions as the original problem.

For the 0^{th} order term it is well known that this homogeneous ODE has a set of solutions (normalized sine functions) for $\theta^{(0)}$ and corresponding eigenvalues for $\lambda^{(0)}$.

We then take the inner products, $\theta^{(1)}$ with the 0^{th} order equation and $\theta^{(0)}$ with the first order equation subtracting we get the expression

$$\lambda^{(1)} \int_0^h \theta^{(0)}(s) ds = \lambda^{(1)} - \int_0^h (\lambda^{(0)} - k^2) \phi^{(1)}(0) \theta^{(0)}(s) ds ;$$

which may be solved if we know the value of the constant $\phi^{(1)}(0)$.



THE TRANSMUTATION RESULTS

THE CORRECTIONS TO FIRST ORDER IN ϵ ARE

$$\lambda_n = \lambda_n^{(0)} + \epsilon \left[\frac{(2n-1)\pi}{2h} \sqrt{\frac{2}{h}} \phi_n^{(1)}(0) \right]$$

AND

$$\begin{aligned} \psi_n(z) = \theta_n^{(0)}(z) + \epsilon \left[\phi_n^{(1)}(0) \left(\left| 1 - \frac{z}{h} \right| \cos \frac{(2n-1)\pi z}{2h} - \int_0^h \left| 1 - \frac{z}{h} \right| \cos \frac{(2n-1)\pi z}{2h} \theta_n^{(0)}(z) dz \right) \right. \\ \left. + \left(\int_h^z K^{(1)}(z,s) \theta_n^{(0)}(s) ds - \int_0^h \left| \int_h^z K^{(1)}(z,s) \theta_n^{(0)}(s) ds \right| \theta_n^{(0)}(z) dz \right) \right]. \end{aligned}$$

Graph 16

We can express the perturbed eigenvalues to the first order in ϵ as

$$\lambda_n = \lambda_n^{(0)} + \epsilon \left[\frac{(2n-1)\pi}{2h} \sqrt{\frac{2}{h}} \phi_n^{(1)}(0) \right]$$

and the perturbed eigenfunctions as

$$\begin{aligned} \psi_n(z) = \phi_n(z) \\ + \epsilon \left\{ \phi_n^{(1)}(0) \left[\left| 1 - \frac{z}{h} \right| \cos \frac{(2n-1)\pi z}{2h} - \int_0^h \left| 1 - \frac{z}{h} \right| \cos \frac{(2n-1)\pi z}{2h} \theta_n^{(0)}(z) dz \right] \right. \\ \left. + \left[\int_h^z K^{(1)}(z,s) \theta_n^{(0)}(s) ds - \int_0^h \left[\int_h^z K^{(1)}(z,s) \theta_n^{(0)}(s) ds \right] \theta_n^{(0)}(z) dz \right] \right\}. \end{aligned}$$

While the correction to the eigenfunction may look complicated it is the whole correction. We have an explicit formula without having to resort to Fourier series expansion.



THE TRANSMUTATION RESULTS (Cont'd)

THE CORRECTIONS TO FIRST ORDER IN ϵ ARE DEFINED
IN TERMS OF

$$\phi_n^{(1)}(0) = \int_0^h K_n^{(1)}(0,s) \theta_n^{(0)}(s) ds$$

AND

$$\epsilon K^{(1)}(z,s) = -\frac{k^2}{2} \left(\int_h^{\frac{z+s}{2}} [n^2(\zeta) - 1] d\zeta + \int_h^{\frac{z-s+2h}{2}} [n^2(\zeta) - 1] d\zeta \right).$$

GRAPH 17

The corrections for the eigenvalues and eigenfunctions are expressed in terms of $\phi_n^{(1)}(0)$ and $K^{(1)}(z,s)$. The first constant is defined by the expression

$$\phi_n^{(1)}(0) = \int_0^h K_n^{(1)}(0,s) \theta_n^{(0)}(s) ds,$$

and the first order in epsilon term of the power series expansion of the kernel is given by

$$\epsilon K^{(1)}(z,s) = -\frac{k^2}{h} \left[\int_h^{\frac{(z+s)/2}{2}} [n^2(\zeta) - 1] d\zeta + \int_h^{\frac{(z-s)/2}{2} + h} [n^2(\zeta) - 1] d\zeta \right].$$

We again have the perturbation type result where the corrections are explicit functions of θ , the unperturbed eigenfunction, λ , the unperturbed eigenvalue and the expression $[n^2(z) - 1]$ which is the perturbation.



CONSISTENCY OF THE TWO APPROACHES

RECALL

$$\begin{aligned}\epsilon\lambda_n^{(1)} &= \epsilon \left[\frac{(2n-1)\pi}{2h} \sqrt{\frac{2}{h}} \phi_n^{(1)}(0) \right] \\ &= \frac{(2n-1)\pi}{2h} \sqrt{\frac{2}{h}} \int_0^h \frac{k^2}{2} \left(\int_h^{\frac{s}{2}} [n^2(\zeta) - 1] d\zeta + \int_{\frac{s}{2}+h}^s [n^2(\zeta) - 1] d\zeta \right) \phi_n^{(0)}(s) ds.\end{aligned}$$

INTEGRATION BY PARTS GIVES

$$\epsilon\lambda_n^{(1)} = \frac{2k^2}{h} \int_0^h (n^2(\zeta) - 1) \left[\sin \frac{(2n-1)\pi\zeta}{2h} \right]^2 d\zeta.$$

Graph 18

The results of the two methods look dissimilar, however we can show that the results are consistent. To show consistency of the first order correction of the eigenvalues we start with

$$\epsilon\lambda_n^{(1)} = \epsilon \left[\frac{(2n-1)\pi}{2h} \sqrt{\frac{2}{h}} \phi_n^{(1)}(0) \right].$$

We substitute for $\phi_n^{(1)}(0)$, including the power series expression for the kernel function. Integration by parts yields

$$\epsilon\lambda_n^{(1)} = \frac{2k^2}{h} \int_0^h [n^2(\zeta) - 1] \left[\sin \frac{(2n-1)\pi\zeta}{2h} \right]^2 d\zeta,$$

which is identical to Titchmarsh.



WE CAN SHOW THAT THE FOURIER COEFFICIENTS OF THE FIRST ORDER CORRECTION DEFINED BY

$$\alpha_{nm} = \int_0^h \psi_n^{(1)}(z) \theta_m^{(0)}(z) dz,$$

ARE GIVEN BY

$$\epsilon \alpha_{nm} = \frac{-2k^2}{h(\lambda_m^{(0)} - \lambda_n^{(0)})} \int_0^h (n^2(z) - 1) \sin \frac{(2n-1)\pi z}{2h} \sin \frac{(2m-1)\pi z}{2h} dz$$

AND

$$\epsilon \alpha_{nn} = 0.$$

Graph 19

The correction given by the transmutation is radically different from the classical results. However if we express the transmutation correction in Fourier series with respect to the functions $\theta^{(0)}$, we see that the Fourier coefficients

$$\alpha_{nm} = \int_0^h \psi_n^{(1)}(z) \theta_m^{(0)}(z) dz,$$

result.

After again making substitutions for $\phi^{(1)}(0)$ and $K^{(1)}(z,s)$, several integrations by parts and some changes of variable these coefficients may be used to derive the classical Fourier coefficients

$$\epsilon \alpha_{nm} = \frac{-2k^2}{h(\lambda_m^{(0)} - \lambda_n^{(0)})} \int_0^h [n^2(z) - 1] \sin \frac{(2n-1)\pi z}{2h} \sin \frac{(2m-1)\pi z}{2h} dz$$

and

$$\epsilon \alpha_{nn} = 0.$$

Abraham, Phil
Code 5132, Physical Acoustic Branch
Naval Research Lab.
Washington, DC 20375

Ahluwalia, D. S.
Chairman
Mathematics Department
New Jersey Institute of Technology
Newark NJ 07102

Ames, W. F.
Georgia Inst of Tech
Atlanta, GA 30332

Andrushkiw, R.
Mathematics Department
New Jersey Institute of Technology
Newark NJ 07102

Avula, Xavier
Institute for Applied Science
Box 1488
Rolla, MI 65401

Baer, Ralph
Naval Research Lab.
Washington, DC 20375

Baggeroer, Authur B.
Mass Inst of Tech
Cambridge, MA 02139

Bates, Susan
Syscon
Middletown, RI 02840

Bayliss, Alvin
Exxon, Res
P.O. Box 45
Linden, NJ 07036

Bleistein, Norman
Mathematics Department
Colorado School of Mines
Golden, CO 80401

Bojarski, Norbert
16 Pine Valley Lane
Newport Beach, CA 92660

Bragg, Louis
Department of Mathematics
Oakland University
Rochester, MI 48063

Burridge, Robert
Courant Institute of Mathematical Sciences
251 Mercer St
New York, NY 10012

Carroll, Robert
Department of Mathematics
University of Ill
Urbana, IL 61801

Cohn, Harvey
Mathematics Department
Convent Ave and 138th St
New York, NY 10031

Caviness, B. F.
Professor and Chairman
Department of Computer and
Information Sciences
University of Delaware
Newark, DE 19716

Colton, David
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Davis, J.
Planning Systems, Inc.
Suite B, Foss Building
95 Trumbull Street
New London, CT 06320

Yves Desaubies
Woods Hole Oceanographic Inst.
Woods Hole, MA 02543

DeSanto, John
Mathematics Department
Colorado School of Mines
Golden, CO 80401

Driver, R.
Mathematics Department
University of Rhode Island
Kingston, RI 02881

Dozier, Lewis
Sciences Applications, Inc.
1710 Goodridge Dr.
McLean, VA 22102

Dout, J.
Woods Hole Oceanographic Inst.
Woods Hole, MA 02543

Ebert, G.
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Elman, Howard
Computer Sciences Department
University of Maryland, Baltimore County
Cantonville, MD 21228

Evans, Richard
ODSI Defense Systems, Inc.
N. Stonington Prof. Center
N. Stonington CT 06359-1738

Felsen, L. B.
Dept of Elec Eng and
Computer Science
Polytechnic University
Farmingdale, NY 11735

Finizio, N.
Mathematics Department
University of Rhode Island
Kingston, RI 02881

Fiduccia, Charles
GE, Research and Development
P.O. Box 8
Schenectady, NY 12301

Fishman, Louis
Department of Civil Engineering
The Catholic University of America
Washington, DC 20064

Fleury, Patrick
4801 Shaboygan Ave. Apt-101
Madison, Wisconsin 53705

Frankenthal, Shimshon
Department of Ocean Engineering
MIT
Cambridge, MA 02139

Gilbert, Robert P.
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Goodman, Ralph
SACLANT ASW Research Center
APO New York 09019

Grove, E.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Hewer, Gary
Naval Weapons Center
China Lake, CA 93555

Jacobson, Melvin
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, NY 12181

Jeffrey, Allen
Department of Engineering Mathematics
University of Newcastle Upon Tyne
NE 171U
England

Johnson, D.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Kleinman, Ralph
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Kriegsmann, Gregory A.
Dept of Eng Sci and Appl Math
Northwestern University
Evanston, IL 60201

Kot, Alex
Electrical Engineering
University of Rhode Island
Kingston, RI 02881

Kuo, John T.
Aldridge Lab of Appl Geophysics
Columbia Univ
New York, NY 10027

Ladas, G.
Mathematics Department
University of Rhode Island
Kingston, RI 02881

Lakshmanasamy, S.
Electrical Engineering
University of Rhode Island
Kingston, RI 02881

Lamagna, Edmund
Computer Sciences Department
University of Rhode Island
Kingston, RI 02881

Lengyel, Grabriel
Department of Electircal Engineering
University of Rhode Island
Kingston, RI 02881

Lewis, J.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Li, Dong-Jye
Ocean Engineering Department
University of Rhode Island
Kingston, RI 02881

Li, Jian-Shu
Box 2155, Yale Station
New Haven, CT 06520

Liu, P.
Mathematics Department
University of Rhode Island
Kingston, RI 02881

Lu, I. T.
Dept of Elec Eng and
Computer Science
Polytechnic University
Farmingdale, NY 11735

Madych, Wolodymyr
Department of Mathematics
University of Connecticut
Storrs, CT 06268

McCoy, J. J.
Department of Mathematics
Catholic University of America
Washington, DC 20064

Maltz, Frederick
Westinghouse Electric Corp.
P.O. Box 1488 - M/S 9260
Anapolis, MD 21404

Marshall, Charles
State University of New York
Computer Science Dept.
Potdam, NY 13676

Mayfield, Betty
22 E 4th Street
Frederick, MD 21701

Merklinger, Harold
Defense Research Establishment Atlantic
9 Grove Street
Darthmouth, Nova Scotia
Canada

Miller, Willard Jr.
School of Mathematics
University of Minnesota
Minneapolis, MN 55455

Montgomery, J.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Mullikin, Thomas
Code 411MA
Office Naval Research
800 Quincy Street
Arlington, VA 22217

Norris, A. N.
Department Computer Sci.
Rutgers Univ
New Brunswick, NJ 08903

Pakula, Lewis
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Papadakis, John
Department of Mathematics
University of Crete
Iraklion
Greece

Pierce, Allan D.
Georgia Inst of Tech
Atlanta, GA 30332

Porter, Michael
Code 5120
Naval Research Lab.
Washington, DC 20375

Richards, P. C.
IBM Corporation
Atlanta, GA 30332

Reiss, Edward L.
Dept of Eng Sci and Appl Math
Northwestern University
Evanston, IL 60201

Robertson, John
Department of Mathematics
U.S. Military Academy
West Point, NY 10996-1786

Rokhlin, Vladimir
Computer Science Dept.
Yale University
10 Hillhouse
New Haven, CT 06520

Roxin, E.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Shisha, O.
Mathematics Department
University of Rhode Island
Kingston, RI 02881

Sine, Robert
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Santosa, Fadil
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Sarkissian, Angie
Sach/Freeman Associates
Bowie, MD 20715

Saunders, D.
Department of Computer and
Information Sciences
University of Delaware
Newark, DE 19716

Schmidt, Henrik
SACLANT ASW Research Centre
APO New York 09019

Schultz, Martin
Computer Science Department
10 Hillhouse
New Haven, CT 06520

Schwartz, Stuart C.
Department of Elec. Engineering
Princeton University
Princeton, NJ 08544

Shen, Longjun
P.O. Box 8009
Beijing
People's Republic of China

Siegmann, Bill
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, NY 12181

Spofford, Charles
Sciences Applications, Inc.
1710 Goodridge Dr.
McLean, VA 22102

St. Mary, Donald
Cntr for Appl Math & Sci Comp
Department of Mathematics
University of Mass
Amherst, MA 01003

Stakgold, Ivar
Chairman
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716

Stepanishen, Peter
Department of Ocean Engineering
University of Rhode Island
Kingston, RI 02881

Sternberg, Robert
Office Naval Research
495 Summer Street
Boston, MA 02210

Stickler, David
Mathematics Department
Colorado School of Mines
Golden, CO 80401

Suryanaryan, E.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Teng, Yu-chiung
Aldridge Lab of Appl Geophysics
Columbia Univ
New York, NY 10027

Thomson, David J.
Defence Research Establishment Pacific
Forces Mail Office
Victoria, British Columbia
Canada VOS 1B0

Trefethen, Lloyd N.
Department of Mathematics
Mass Inst of Tech
Cambridge, MA 02139

Tufts, Donald
Department of Electrical Engineering
University of Rhode Island
Kingston, RI 02881

Turkel, Eli
Tel Aviv University
Math Science School
Ramat Aviv
Tel Aviv, Israel

Van Loan, Charles
Department of Computer Science
Cornell University
Ithaca, NY 14853

Verma, Ghasi R.
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

Vichnevetsky, R
Department Computer Sci.
Rutgers Univ
New Brunswick, NJ 08903

Vignes, J.
Universite P. et M. Curie
4, Place Jussieu
75230 PARIS Cedex
France

Weinberg, H.
Senior Technical Director
ODSI Defense Systems, Inc.
N. Stonington Prof. Center
N. Stonington CT 06359-1738